

Literatuurlijst SOP Remifentanil PCA als methode van pijnstilling tijdens de baring.

Richtlijn Medicamenteuze pijnbehandeling tijdens de bevalling.
CBO richtlijn 2008.

Glass PS, Hardman D, Kamuyama Y et al.
Preliminary pharmacokinetics and pharmacodynamics of an ultra-short acting opioid.
Anesthesia and Analgesia 1993 77: 1031-40.

Kan RW, Hughes SC, Rosen M.A, Kossin C, Preston PG, Lobo EP.
Intravenous remifentanil: placental transfer, maternal and neonatal effects.
Anesthesiology 1998 88: 1467-74.

Hinova A, Fernando R.
Systemic remifentanil for labor analgesia.
Anesthesia and Analgesia 2009 109: 1925-9.

Volmanen P, Akural EC, Raudaskoski T, Alahuta S.
Remifentanil in obstetric analgesia: a dose-finding study.
Anesthesia and Analgesia 2002 94: 913-7.

Blair JM, Dobson GT, Hill DA, Mc Cracken GR, Fee JP.
Patient-controlled analgesia for labour: a comparison of remifentanil with pethidine.
Anaesthesia 2005 60:22-7.

Volikas I, Butwick A, Wilkinson C, Fleming A, Nicholson G.
Maternal and neonatal side-effects of remifentanil patient-controlled analgesia in labour.
British Journal of Anaesthesia 2005 95: 504-9.

Babenco HD, Conard PF, Gross J.B.
The pharmacodynamic effect of a remifentanil bolus on ventilatory control.
Anesthesiology 2000 92: 393-8.

Harbers J, Drogdrop A, van Ieperen R.
Remifentanil is safe and effective for patient-controlled analgesia or labour.
International Journal of Obstetric Anaesthesia 2008 17:s10

Douma MR, Verwey RA, Kam-Endtz CE, van der Linden PD, Stienstra R.
Obstetric analgesia: a comparison of patient-controlled meperidine, remifentanil and fentanyl in labour.
British Journal of Anaesthesia 2010 104: 209-

Melber AA, Reinhardt D, Bansi AS.
The launch of remifentanil PCA in labour in Switzerland combined with web-based continuous quality control.
International Journal of Obstetric Anaesthesia 2011 20:s35

Schnabel A, Hahn N, Brosheit J et al.

Remifentanil for labour analgesia: a meta-analysis of randomised controlled trials.
European Journal of Anaesthesiology 2012 29: 177-85

Hill D.

Remifentanil patient-controlled analgesia should be routinely available for use in labour- Pro

International Journal of Obstetric Anaesthesia 2008 17: 336-9

Van de Velde M.

Remifentanil patient-controlled analgesia should be routinely available for use in labour- Con

International Journal of Obstetric Anaesthesia 2008 17: 339-42

Huchatuta NA, Kinsella SM.

Remifentanil for labour analgesia: time to draw a breath?

Anaesthesia 2013 68: 227-35

Marr R, Hiyams J, Bythell V.

Cardiac arrest in an obstetric patient using remifentanil patient-controlled analgesia.

Anaesthesia 2013 68

Pruefer C, Bewlay A.

Respiratory arrest with remifentanil patient-controlled analgesia-another case.

Anaesthesia 2012 67: 1044-5.

Bonner JC, McClymont W.

Respiratory arrest in an obstetric patient using remifentanil patient-controlled analgesia.

Anaesthesia 2012 67: 538-40.

Kranke P et al.

Must we press on until a young mother dies? Remifentanil patient-controlled analgesia in labour may not be suited as a "Poor man's epidural"

B.M.C. Pregnancy and Childbirth 2013 13: 139

O'Sullivan G, Liu B, Hart D, Seed P, Shennan A.

Effect of food intake during labour on obstetric outcome: randomised controlled trial.

British Medical Journal 2009 338: b784

Douma MR, Middeldorp JM, Verwey RA, Stienstra R.

A randomised comparison of intravenous remifentanil patient-controlled analgesia with epidural ropivacaine/sufentanil during labour.

International Journal of Obstetric Anaesthesia 2011 20: 118-23

Freeman LM, Bloemenkamp KW, Franssen MT, Papatsonis DN, Hajenius PJ, van Huizen ME, Bremer HA, van den Akker ES, Woiski MD, Porath MM, van Beek E, Schuitemaker N, van der Salm PC, Fong BF, Radder C, Bax CJ, Sikkema M, van den Akker-van Marle ME, van Lith JM, Lopriore E, Uildriks RJ, Struys MM, Mol BW, Dahan A, Middeldorp JM.

Remifentanil patient-controlled analgesia versus epidural analgesia in labour. A multi-center randomized controlled trial.
BMC Pregnancy Childbirth 2012 12:63

Schuurhuis A, Roumen FJ, de Boer JB; werkgroep Anesthesiologie.
Practice guideline 'Pharmaceutical pain treatment during labour'; the woman's request is sufficient Indication.
Nederlands Tijdschrift voor Geneeskunde 2009 153:A551.

Birnbach D, Ranasinghe S.
Is remifentanil a safe and effective alternative to neuraxial labor analgesia? It all depends.
Anesthesia and Analgesia 2014 118: 491-3.

Stocki D, Matot I, Einav S, Ginosar Y, Eventov-Friedman S, Weiniger CF.
Randomised controlled trial of Efficacy and Respiratory effects of patient-controlled Intravenous Remifentanil analgesia and patient-controlled Epidural analgesia in laboring women.
Anesthesia and Analgesia 2014 118 589-97.